

09/913,325

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 August 2000 (31.08.2000)

PCT

(10) International Publication Number
WO 00/049937 A3

(51) International Patent Classification⁷: C07H 21/04,
A61K 48/00, C12N 15/85, C12Q 1/68

(21) International Application Number: PCT/US00/04875

(22) International Filing Date: 25 February 2000 (25.02.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/121,726 26 February 1999 (26.02.1999) US

(71) Applicant (*for all designated States except US*): THE
UNIVERSITY OF BRITISH COLUMBIA [CA/CA];
2075 Westbrook Mall, Vancouver, British Columbia V6T
1Z1 (CA).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): GLEAVE, Martin
[CA/CA]; 4693 Drummond Drive, Vancouver, British Co-
lumbia V6R 1E8 (CA). RENNIE, Paul, S. [CA/CA]; 3731
Ullsmore Avenue, Richmond, British Columbia V7C 1S3
(CA). MIYAKE, Hideaki [JP/CA]; 2638 Ash Street, #204,
Vancouver, British Columbia V5Z 4K3 (CA). NELSON,
Colleen [US/CA]; 3299, 137A Street, Surrey, British Co-
lumbia V4P 2B5 (CA).

(74) Agents: PARSONS, Nancy, J. et al.; Oppedahl & Larson
LLP, P.O. Box 5068, Dillon, CO 80435-5068 (US).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,
UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:
— with international search report

(88) Date of publication of the international search report:
7 December 2000

(48) Date of publication of this corrected version:
11 July 2002

(15) Information about Correction:
see PCT Gazette No. 28/2002 of 11 July 2002, Section II

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: TRPM-2 ANTISENSE THERAPY

(57) Abstract: It has now been determined that antisense therapy which reduces the expression of TRPM-2 provides therapeutic benefits in the treatment of cancer. In particular, such antisense therapy can be applied in treatment of prostate cancer and renal cell cancer. Addition of antisense TRPM-2 ODN to prostatic tumor cells *in vivo* is effective for delaying the onset of androgen inde-
pendence. Thus, prostate cancer can be treated in an individual suffering from prostate cancer by initiating androgen-withdrawal
to induce apoptotic cell death of prostatic tumor cells in the individual, and administering to the individual a composition effective
to inhibit expression of TRPM-2 by the tumor cells, thereby delaying the progression of prostatic tumor cells to an androgen-inde-
pendent state in an individual. Combined use of antisense TRPM-2 and taxanes synergistically enhances cytotoxic chemosensitivity
of androgen-independent prostate cancer. In addition, it has also been found that antisense TRPM-2 has beneficial effect for other
cancer types. Specifically, antisense TRPM-2 ODN enhances chemosensitivity in human Renal cell cancer, a normally chemore-
sistant disease with no active chemotherapeutic agent having an objective response rate higher than 10 %. Radiation sensitivity is
also enhanced when cells expressing TRPM-2 are treated with antisense TRPM-2 ODN. Thus, the antisense TRPM-2 ODNs can be
used to enhance hormone sensitivity, chemosensitivity and radiation sensitivity of a variety of cancer types in which expression of
TRPM-2 has been observed.



WO 00/049937 A3

- 1 -

TRPM-2 ANTISENSE THERAPY

This application claims priority from US Provisional Patent Application No. 60/121,726, filed February 26, 1999, which is incorporated herein by reference.

Background of the Invention

This application relates to antisense treatments for cancer making use of an antisense oligonucleotide that binds to testosterone-repressed prostate message-2 (TRPM-2).

Prostate cancer is the most common cancer that affects men, and the second leading cause of cancer deaths in men in the Western world. Because prostate cancer is an androgen-sensitive tumor, androgen withdrawal, for example via castration, is utilized in some therapeutic regimens for patients with advanced prostate cancer. Androgen withdrawal leads to extensive apoptosis in the prostate tumor, and hence to a regression of the disease. However, castration-induced apoptosis is not complete, and a progression of surviving tumor cells to androgen-independence ultimately occurs. This progression is the main obstacle to improving survival and quality of life, and efforts have therefore been made to target androgen-independent cells. These efforts have focused on non-hormonal therapies targeted against androgen-independent tumor cells (Yagoda et al., *Cancer* 71 (Supp. 3): 1098-1109 (1993); Oh et al., *J. Urol.* 60: 1220-1229 (1998)), however, so far no non-hormonal agent has improved survival. Alternative approaches are therefore indicated.

It has been observed that numerous proteins are expressed in increased amounts by prostate tumor cells following androgen withdrawal. At least some of these proteins are assumed to be associated with the observed apoptotic cell death which is observed upon androgen withdrawal. (Raffo et al., *Cancer Res.*: 4448-4445 (1995); Krajewska et al., *Am. J. Pathol.* 148: 1567-1576 (1996); McDonnell et al., *Cancer Res.* 52: 6940-6944 (1992)). The functions of many of the proteins, however, is not clearly understood. TRPM-2 (also known as sulfated glycoprotein-2 (SGP-2) or clusterin) is within this latter category.

TRPM-2 is a ubiquitous protein, with a diverse range of proposed activities. In prostate epithelial cell, expression of TRPM-2 increases immediately following castration, reaching peak levels in rat prostate cells at 3 to 4 days post castration, coincident with the onset of massive

- 2 -

cell death. These results have led some researchers to the conclusion that TRPM-2 is a marker for cell death, and a promoter of apoptosis. On the other hand, the observation that Sertoli cells and some epithelial cells express high levels of TRPM-2 without increased levels of cell death, raises questions as to whether this conclusion is correct.

Sensibar et al., *Cancer Research* 55: 2431-2437 (1995) reported on *in vitro* experiments performed to more clearly elucidate the role of TRPM-2 in prostatic cell death. They utilized LNCaP cells transfected with a gene encoding TRPM-2 and observed whether expression of this protein altered the effects of tumor necrosis factor α (TNF α), to which LNCaP cells are very sensitive, with cell death normally occurring within about 12 hours. Treatment of the transfected LNCaP cells with TNF α was shown to result in a transient increase in TRPM-2 levels for a period of a few hours, but these levels had dissipated by the time DNA fragmentation preceeding cell death was observed. Using an antisense molecule corresponding to the bases 1-21 of the TRPM-2 sequence, but not other TRPM-2 antisense oligonucleotides, resulted in a substantial reduction in expression of TRPM-2, and an increase in apoptotic cell death in LNCaP cells exposed to TNF α . This led Sensibar et al. to the hypothesis that overexpression of TRPM-2 could protect cells from the cytotoxic effect of TNF, and that TRPM-2 depletion is responsible for the onset of cell death, although the mechanism of action remains unclear.

While Sensibar et al. provides information about the possible role of TRPM-2, it nevertheless discloses results from only a model system in which expression of TRPM-2 is based on a transfected gene. Furthermore, expression levels of TRPM-2 is very low or absent in LNCaP cells grown in other labs. The situation which results *in vivo* when prostate tumor cells are subjected to androgen withdrawal is far more complex, with numerous proteins changing expression levels as a result. Thus, it is not possible from the Sensibar et al. data to predict whether TRPM-2 would perform the same function when present in combination with other proteins, or whether changes in levels of TRPM-2 following androgen withdrawal *in vivo* could provide any therapeutic benefits. Indeed, the fact that TRPM-2 is expressed in substantial quantities in prostatic tumor cells at various stages following androgen withdrawal, including stages where significant apoptotic cell death is occurring suggests that role of TRPM-2 *in vivo* may be more complicated. Thus, while the art provides data concerning certain aspects of apoptotic cell death in prostatic tumor cells, it offers

- 3 -

neither a teaching or a suggestion of a methodology to provide a delay in the onset of androgen-independence.

It is an object of the present invention to provide such a method.

It is a further object of the present invention to provide therapeutic antisense molecules for delaying the onset of androgen independence in prostatic tumor cells.

It is an additional object of the present invention to provide a method for enhancing the chemosensitivity or radiation sensitivity of cancer cells from a cancer that expresses TRPM-2.

It is a further object of the present invention to provide therapeutic antisense molecules for inhibiting expression of TRPM-2.

Summary of the Invention

In accordance with the present invention, it has now been determined that antisense therapy which reduces the expression of TRPM-2 provides therapeutic benefits in the treatment of cancer. In particular, such antisense therapy can be applied in treatment of prostate cancer and renal cell cancer.

Addition of antisense TRPM-2 oligodeoxynucleotide (ODN) to prostatic tumor cells *in vivo* is effective for delaying the onset of androgen independence. Thus, in one aspect, the invention provides a method for treating prostate cancer in an individual suffering from prostate cancer, comprising the steps of initiating androgen-withdrawal to induce apoptotic cell death of prostatic tumor cells in the individual, and administering to the individual a composition effective to inhibit expression of TRPM-2 by the tumor cells, thereby delaying the progression of prostatic tumor cells to an androgen-independent state in an individual. Furthermore, combined use of antisense TRPM-2 plus cytotoxic chemotherapy (e.g. taxanes) synergistically enhances chemosensitivity in hormone refractory prostate cancer. In another aspect of the invention, a second antisense ODN which inhibits expression of an anti-apoptotic protein other than TRPM-2 is administered along with the antisense TRPM-2 ODN.

It has also been found that antisense TRPM-2 has beneficial effects for other cancer types. Specifically, antisense TRPM-2 ODN enhances chemosensitivity in human Renal cell cancer, a normally chemoresistant disease with no active chemotherapeutic agent having an

- 4 -

objective response rate higher than 10%. Radiation sensitivity is also enhanced when cells expressing TRPM-2 are treated with antisense TRPM-2 ODN. Thus, the antisense TRPM-2 ODNs can be used to treat a variety of cancer types in which expression of TRPM-2 has been observed.

Brief Description of the Figures

Fig. 1 shows the delay in onset of androgen-independence which is achieved using an antisense TRPM-2 ODN;

Fig. 2 shows the positions of 10 antisense oligonucleotides evaluated for the ability to inhibit TRPM-2 expression and delay onset of androgen-independence;

Fig. 3 shows expression levels of TRPM-2 mRNA in the presence of various antisense ODNs;

Fig. 4 shows the levels of TRPM-2 mRNA in Shionogi cells treated *in vitro* with varying amounts of antisense TRPM-2 ODN or a mismatch control;

Fig. 5 shows the dose-response curve for combinations of taxol and antisense TRPM-2 ODN;

Fig. 6 shows the dose-response curve for combinations of taxol, antisense TRPM-2 ODN and antisense Bcl-2 ODN;

Fig. 7A shows decrease in TRPM-2 mRNA levels in human renal cell cancer after treatment with antisense TRPM-2 ODNs;

Fig. 7B shows the increase in chemosensitivity of human renal cell cancer to taxol after treatment with antisense TRPM-2 ODNs;

Fig. 8 shows TRPM-2 expression in PC-3 prostate cancer cells after various doses of radiation;

Figs. 9A and 9B show the comparative radiation resistance of human prostate cell lines which overexpress (LNCaP/T) and normally (LNCaP/P) express TRPM-2;

Fig. 10 shows the increased susceptibility of PC-3 cells to radiation after treatment with antisense TRPM-2 ODN; and

- 5 -

Figs. 11A and 11B show the increased sensitivity of PC-3 cells to radiation after treatment with antisense TRPM-2 ODN.

Figs. 12A and 12B show the increased sensitivity of Shionogi tumor cells to chemotherapy agents paclitaxel and mitoxanthrone when administered with antisense TRPM-2 ODN.

Detailed Description of the Invention

The present invention relates to antisense TRPM-2 ODNs and to the use of these compositions in the treatment of cancer. The invention can be applied in the treatment of cancers where the cancer cells express TRPM-2. Three significant classes of cancer cells which express TRPM-2 are prostate cancer cells, human renal cell cancer (RCC) cells and some breast cancer cells.

In one embodiment, the present invention provides a method for enhancing castration-induced tumor cell death and delaying the progression of prostatic tumor cells to androgen independence; a therapeutic method for the treatment of individuals, including humans, suffering from prostate cancer; and therapeutic agents effective for use in such methods. The therapeutic method of the invention will most commonly be used in the treatment of individuals with advanced prostate cancer.

Enhancement of castration-induced tumor cell death and delay of the progression of androgen-sensitive prostatic cancer cells to androgen-independent is achieved by inhibiting the expression of TRPM-2 by the cells. Experiments were performed in three model systems, the *in vivo* Shionogi tumor model, the human TRPM-2 transfected LNCaP model, and the human PC-3 model, which taken together demonstrated that such inhibition leading to delay of androgen-independence can be achieved by treating androgen-sensitive prostatic tumor cells with antisense oligodeoxynucleotides (ODNs).

In the first experiment, the ability of a mouse TRPM-2 antisense molecule, (Seq. ID. No. 1) to delay onset of androgen independence in the Shionogi tumor model was evaluated. The Shionogi tumor model is a xenograft of an androgen-dependent mouse mammary carcinoma that grows subcutaneously in male syngeneic hosts. Shionogi tumor cells are highly tumorigenic and

- 6 -

locally invasive. The cells have been shown to respond to androgen withdrawal in a manner which mimics the observed behavior of prostatic tumor cells, and have been accepted as a valid model for prostate cancer in humans. (Bruchovsky et al., *Cancer Res.* 50: 2275-2282 (1990); Rennie et al., *Cancer Res.* 48: 6309-6312 (1988); Bruchovsky et al., *Cell* 13: 272-280 (1978); Gleave et al., in *Genitourinary Oncology*, pp. 367-378, Lange et al., eds, Lippencott (1997); Gleave et al., *J. Urol.* 157: 1727-1730 (1997); Bruchovsky et al., *The Prostate* 6: 13-21 (1996)). Thus, androgen withdrawal precipitates apoptosis and tumor regression in a highly reproducible manner. Further, changes in expression of TRPM-2 and Bcl-2 in human prostate cancer following castration and during progression to androgen independence are similar to those observed in Shionogi tumor cells. Thus, the Shionogi tumor model mimics many of the characteristics of prostate cancer cells. Further, the Shionogi tumor model provides a very useful model for the evaluation of the ability of compounds to delay the onset of androgen-independence. Despite complete tumor regression after castration, rapidly growing androgen-independent Shionogi tumors invariably recur after one month, which provides a reliable end point to evaluate agents which can delay the progression to androgen-independence. In general, events which occur in the Shionogi tumor model within one month occur in human patients within about two years.

The ability of the antisense ODNs that inhibit expression of TRPM-2 to delay the onset of androgen-independence was evaluated by measuring tumor volume post-castration in the Shionogi tumor model. The test animals (n=7) were treated intraperitoneally once daily with 12.5 mg/kg repeat doses of antisense TRPM-2 ODNs (Seq. ID. No 1) in a buffered saline solution. As a control, animals (n=7) were treated with a mismatch ODN (Seq. ID. No. 2). As shown in Fig. 1, both test and control groups showed the expected decline in tumor volume immediately following castration, but the tumors in the antisense TRPM-2 ODN-treated mice regressed faster than the controls. The control group also exhibited the expected increase in tumor volume which is associated the development of androgen-independence. In contrast, at 49 days post-castration, little tumor regrowth had occurred in the mice treated using the antisense TRPM-2 ODN. Tumors did eventually recur in the antisense TRPM-2 ODN-treated mice, but the median time to recurrence is approximately twice that of the control group. Thus, inhibition of TRPM-2 is effective not only for increasing the amount of cell death which occurs immediately following androgen withdrawal,

- 7 -

but also for delaying the onset of androgen-independence. The more rapid decrease in tumor volume in the mice treated with antisense TRPM-2 ODNs was due to earlier onset and more extensive castration-induced apoptosis. This was confirmed by detecting poly(ADP-ribose) polymerase (PARP) cleavage fragments in Shionogi tumor specimens (Miyake, et al., *Cancer Res.* 60:170-176 (2000)).

To evaluate which human antisense ODNs complementary to TRPM-2 mRNA sequences are most effective for this purpose, a series of ten antisense phosphorothioate ODNs were prepared spanning various mRNA regions as shown in Fig. 2. The sequences of these ten ODNs are set forth in the attached Sequence Listing as Seq. ID. Nos. 3-12. The ten human antisense ODNs were evaluated using TRPM-2 transfected LNCaP cells and human prostate cancer PC-3 cells for their ability to inhibit expression of TRPM-2 mRNA. As shown in Fig.3, the antisense ODNs tested produced variable levels of inhibition of TRPM-2 mRNA expression, with the best results being achieved with Seq. ID Nos. 4, 5, and 12. Sequence ID No. 5 corresponds to the sequence used by Sensibar et al. that produced inhibition of TRPM-2 expression in LNCaP cells, and is complementary to the first 21 bases of the TRPM-2 mRNA. The most effective down-regulation occurred with Seq. ID No. 4. Common to all of the effective sequences is an overlap with either the initiation or termination sites of the TRPM-2 mRNA. Thus, in a general sense, the method of the invention can be practiced with antisense oligonucleotides which are complementary to a region of the TRPM-2 mRNA spanning either the translation initiation site or the termination site.

In accordance with a further aspect of with the invention, therapeutic treatment of individuals, including human individuals, suffering from prostate cancer is achieved by initiating androgen-withdrawal to induce apoptotic cell death of prostatic tumor cells in the individual, and administering to the individual a composition effective to inhibit expression of TRPM-2 by the tumor cells, thereby delaying the progression of prostatic tumor cells to an androgen-independent state in an individual.

Initiation of androgen withdrawal may be accomplished via surgical (removal of both testicles) or medical (drug-induced suppression of testosterone) castration, which is currently indicated for treatment of prostate cancer. Medical castration can be achieved by various regimens,

- 8 -

including LHRH agents or antiandrogens. (Gleave et al., *CMAJ* 160: 225-232 (1999)).

Intermittent therapy in which reversible androgen withdrawal is effected is described in Gleave et al. *Eur. Urol.* 34 (Supp. 3): 37-41 (1998).

The inhibition of TRPM-2 expression may be transient, and ideally should occur coincident with androgen withdrawal. In humans, this means that inhibition of expression should be effective starting within a day or two of androgen withdrawal and extending for about 3 to 6 months. This may require multiple doses to accomplish. It will be appreciated, however, that the period of time may be more prolonged, starting before castration and extending for substantial time afterwards without departing from the scope of the invention.

Antisense TRPM-2 ODNs have also been determined to enhance chemosensitivity in human renal cell cancer (RCC). RCC is a chemoresistant disease with no active chemotherapeutic agent with objective response rates higher than 10 %. Increased TRPM-2 expression in renal proximal convoluted cells undergoing apoptosis has been observed after various stimuli including ureteral obstruction and aminoglycosides. However, functional significance of TRPM-2 expression in RCC has not been well documented. Test results show, however, that antisense TRPM-2 ODN enhances chemosensitivity in human RCC CaKi-2 cells (See Example 6, *infra*).

Antisense TRPM-2 ODNs were also found to increase sensitivity to radiation (See Example 7 and Fig. 8).

Inhibition of expression of TRPM-2 may be accomplished by the administration of antisense ODNs, particularly antisense ODNs which are complementary to a region of the TRPM-2 mRNA spanning either the translation initiation site or the termination site. For treatment of prostate cancer in humans, specific useful sequences are those shown in Seq. ID Nos. 4, 5 and 12.

The ODNs employed may be modified to increase the stability of the ODN *in vivo*. For example, the ODNs may be employed as phosphorothioate derivatives (replacement of a non-bridging phosphoryl oxygen atoms with a sulfur atom) which have increased resistance to nuclease digestion. MOE modification (ISIS backbone) is also effective.

Administration of antisense ODNs can be carried out using the various mechanisms known in the art, including naked administration and administration in pharmaceutically acceptable

- 9 -

lipid carriers. For example, lipid carriers for antisense delivery are disclosed in US Patents No. 5,855,911 and 5,417,978 which are incorporated herein by reference. In general, the antisense is administered by intravenous, intraperitoneal, subcutaneous or oral routes, or direct local tumor injection. From the experiments performed using the Shionogi mouse model, it appears that the antisense ODN is preferentially active in the tumor cells. Indeed, TRPM-2 expression in non-tumor tissues was substantially unaffected, and no side effects of the antisense ODN administration were observed.

The amount of antisense ODN administered is one effective to inhibit the expression of TRPM-2 in prostatic cells. It will be appreciated that this amount will vary both with the effectiveness of the antisense ODN employed, and with the nature of any carrier used. The determination of appropriate amounts for any given composition is within the skill in the art, through standard series of tests designed to assess appropriate therapeutic levels.

The method for treating prostate cancer in accordance with the invention may further include administration of chemotherapy agents and/or additional antisense ODNs directed at different targets. For example, it has been found using the Shionogi tumor model that antisense TRPM-2 ODN increases sensitivity to conventional chemotherapy agents such as taxanes (paclitaxel or docetaxel) and mitoxanthrone (Figs. 12A and 12B). As shown in Figs. 12A and 12B, treatment with antisense TRPM-2 ODN in the presence of taxol or mitoxanthrone resulted in a reduced tumor volume as compared to the combination of taxol or mitoxanthrone with the mismatch (MM) ODN. Other agents likely to show synergistic activity include other cytotoxic agents (e.g. cyclophosphamide, topoisomerase inhibitors), angiogenesis inhibitors, differentiation agents and signal transduction inhibitors. Similarly, combinations of TRPM-2 antisense with other antisense species such as antisense Bcl-2 ODN worked better at killing Shionogi cells *in vitro* than either ODN alone. Thus, TRPM-2 can work in concert with other antisense molecules, such as antisense Bcl-2, Bcl-xl and c-myc ODN to provide greater effectiveness.

The invention will now be further described with reference to the following, non-limiting examples.

- 10 -

EXAMPLE 1

Shionogi tumor model experiments were performed using cells from the Toronto subline of transplantable SC-115 AD mouse mammary carcinoma. For *in vivo* studies, approximately 5×10^6 cells of the Shionogi carcinoma were injected subcutaneously in adult male DD/S strain mice. When the Shionogi tumors became 1 to 2 cm in diameter, usually 2 to 3 week after injection, castration was performed through an abdominal incision under methoxyflurane anesthesia. Details of the maintenance of mice, tumor stock and operative procedures have been previously described. Bruchovsky et al., *Cancer res.* 50: 2275-2282 (1990); Rennie et al., *Cancer Res.* 48: 6309-6312 (1988); Bruchovsky et al., *Cell* 13: 272-280 (1978); Gleave et al., in *Genitourinary Oncology*, pp. 367-378, Lange et al., eds, Lippencott (1997); Gleave et al., *J. Urol.* 157: 1727-1730 (1997); Bruchovsky et al., *The Prostate* 6: 13-21 (1996)).

Mice were randomly selected for treatment with murine phosphorothioate antisense TRPM-2 ODN (Seq. ID No. 1) or a mismatch control (Seq. ID No. 2) which is two bases different in sequence from the antisense TRPM-2 ODN. Each experimental group consisted of 7 mice. One day after castration, 12.5 mg/kg of antisense TRPM-2 or mismatch control ODN dissolved in phosphate buffered saline were injected intraperitoneally once daily into each mouse of 40 days. Tumor volume was measured twice weekly, and calculated by the formula length X width X depth X 0.5236. Gleave et al., *Cancer Res.* 52: 1598-1605 (1992). Data points were reported as average tumor volumes \pm standard deviation.

The results of this study are shown in Fig. 1. As shown, Shionogi tumors regressed faster and complete regression occurred earlier in mice treated with antisense TRPM-2 ODN. Furthermore, treatment with antisense TRPM-2 ODN substantially delayed the onset of androgen-independence which is reflected by the increase in tumor volume after day 21 in the control animals. No side effects associated with antisense TRPM-2 or the mismatch control were observed.

To examine the effects of *in vivo* ODN treatment on levels of TRPM-2 mRNA, Northern blot analysis was performed on Shionogi tumor tissue from mice. The mice were treated daily with 12.5 mg/kg of antisense TRPM-2 ODN (n=6) or the mismatch control (n=6) by intraperitoneal injection starting one day after castration. On the fourth day after castration, tumor tissues were harvested and analyzed by Northern blot for TRPM-2 mRNA. Antisense TRPM-2

- 11 -

ODN resulted in a 75% reduction in TRPM-2 mRNA levels in Shionogi tumors compared to mismatch control ODN treated tumors. (Fig. 3).

Comparable analyses were performed on normal mouse organs. Samples of spleen, kidney, prostate and brain were harvested from Shionogi tumor mice treated with antisense TRPM-2 ODN and mismatch control under the same treatment schedule, and analyzed by Northern blot. Although TRPM-2 mRNA levels was significantly lower in tumor tissues, antisense TRPM-2 ODN had no effect on TRPM-2 mRNA levels in the normal organs.

EXAMPLE 2

The sequence selectivity of the antisense TRPM-2 ODN (Seq. ID. No. 1) was confirmed by comparing expression levels of TRPM-2 mRNA in Shionogi tumor cells maintained *in vitro*, after treatment with the varying levels of antisense TRPM-2 ODN or a mismatch control (Seq. ID. No. 2). To facilitate uptake of the ODNs into the cells, the ODNs were formulated in a cationic lipid carrier (Lipofectin™, (Life Technologies, Inc.)). Cells were treated twice over a period of two days using the following protocol. Cells were preincubated for 20 minutes with 4 µg/ml of lipofectin in serum free OPTI-MEM™ (Life Technologies, Inc.) and then incubated with the medium containing the selected concentration of ODN and lipofectin for four hours. The medium was then replaced with the standard culture medium.

The amount of TRPM-2 mRNA in the cells was evaluated using Northern blot analysis. As shown in Fig. 4, treatment of Shionogi cells with antisense TRPM-2 ODN reduced TRPM-2 mRNA levels in a dose dependent manner. In contrast, TRPM-2 mRNA levels were not affected by the mismatch ODN (Seq. ID. No. 2) at any of the employed concentrations. Thus, the affect of antisense TRPM-2 ODN is apparently sequence specific.

EXAMPLE 3

Shionogi cells maintained *in vitro* were treated with varying amounts of taxol alone or in combination with 500 nM antisense TRPM-2 ODN (Seq. ID. No. 1) or the mismatch control (Seq. ID. No. 2). The cells were treated twice, as described in Example 2, and the percentage of viable cells remaining was determined. The results are summarized in Fig. 5. As shown, the

- 12 -

inclusion of antisense TRPM-2 ODN shifted the dose-response curve to the left, lowering the IC_{50} by a factor of 5 to 10. Similar results were achieved using mitoxanthrone in place of paclitaxel (Figs. 12A and 12B).

EXAMPLE 4

The experiment of Example 3 was repeated, with the addition of antisense Bcl-2 ODN (Seq. ID. No. 13) or a mismatch Bcl-2 ODN (Seq. ID. No. 14) in various combinations with antisense/mismatch TRPM-2 ODN and taxol. The results are shown in Fig. 6. The combination of antisense TRPM-2 ODN with antisense Bcl-2 ODN and taxol further enhanced the cytotoxic effects of taxol. Thus, the targeting of additional anti-apoptotic agents appears to provide therapeutic benefits.

EXAMPLE 5

To identify appropriate antisense TRPM-2 ODN sequences for use in human therapy, antisense ODN sequences directed against 10 different sites of the human TRPM-2 gene (Fig. 2, Seq. ID Nos. 3-12) were synthesized and tested for their ability to decrease TRPM-2 gene expression in human prostate cancer PC-3 and transfected LNCaP cells that overexpress TRPM-2 using the same treatment protocol described in Example 2. The results are summarized in Fig. 3. As shown, sequences 4, 5 and 12 are active for reduction of TRPM-2 expression. These three sequences overlap or are immediately adjacent to the translation initiation or termination sites.

EXAMPLE 6

Immunohistochemical staining was used to characterize clusterin expression in 17 RCC and normal kidney tissues obtained from radical nephrectomy specimens. TRPM-2 expression in human renal cancer cell lines ACHN, CaKi-1 and CaKi-2 was evaluated by Northern and Western blot analyses. Northern blot analysis was used to assess changes in TRPM-2 mRNA expression after antisense TRPM-2 ODN treatment. The effects of combined antisense TRPM-2 ODN and taxol treatment on CaKi-2 cell growth was examined using a MTT assay.

- 13 -

Immunostaining showed an increased clusterin expression in 11 RCC specimens in comparison to the adjacent normal kidney tissue. In the remaining 6 cases, no difference was seen between malignant and normal tissue. Both TRPM-2 mRNA and protein expression were detectable in all three human RCC cell lines, with highest levels for CaKi-2.

Antisense TRPM-2 ODN (Seq. ID. No. 1), but not mismatch control ODN (Seq. ID. No. 2), inhibited TRPM-2 expression in CaKi-2 cells in a dose dependant and sequence specific manner (Fig. 7A). Furthermore, antisense TRPM-2 ODN substantially enhanced taxol chemosensitivity, reducing IC₅₀ of taxol by 1 log (500 nM to 50 nM) compared to mismatch control ODN (Fig. 7B). These data demonstrate that TRPM-2 and its protein, clusterin, are expressed at higher levels in RCC compared to normal kidney tissue, and that antisense TRPM-2 ODN may be useful in enhancing the cytotoxic effects of conventional chemotherapy in advanced RCC.

EXAMPLE 7

Antisense TRPM-2 ODNs enhance radiation sensitivity of cancer cells which express TRPM-2. Using northern analysis, we found that radiation therapy results in dose and time dependent increases in TRPM2 gene expression in human prostate cancer PC-3 cells (Fig 8). Overexpression of TRPM2 results in increased resistance to radiation induced cell death. Human prostate LNCaP cells that overexpress TRPM2 (LNCaP/T1) are more resistant to radiation therapy (Figs. 9A and B). Treatment of human prostate cancer PC-3 cells with 100 and 500 nM antisense TRPM-2 ODNs (Seq. ID. NO. 1) significantly reduces cell survival after a single treatment of 4 Gy radiation therapy compared to mismatch ODN (Seq. ID No. 2) treatment. (Fig. 10). Figure 11A and B show dose dependent radiation sensitization of human prostate cancer PC-3 cells after treatment with 10, 50, and 100 nM antisense TRPM-2 oligo *in vitro*.

EXAMPLE 8

To determine whether treatment with human antisense TRPM-2 ODN enhances chemosensitivity in the PC3 human prostate cancer cell line, mice bearing PC3 tumors were treated with antisense human TRPM-2 ODN plus micellar paclitaxel or mitoxantrone, and mismatch control ODN plus micellar paclitaxel or mitoxantrone (Figs. 12A and 12B). ODN was administered for 28

- 14 -

days and either 0.5 mg micellar taxol or 0.3 mg mitoxantrone were administered on two occasions: from day 10 to 14, and day 24 to 28. A significant reduction in tumor size was observed in the antisense ODN treated animals as compared to those treated with mismatch control ODN. This effect was even more pronounced after the second dosing of the micellar paclitaxel or mitoxantrone.

- 15 -

CLAIMS

1. A method for delaying progression of prostatic tumor cells to an androgen- independent state, comprising treating androgen-sensitive prostatic tumor cells with an antisense oligonucleotide which inhibits expression of TRPM-2 by the tumor cells.
2. The method of claim 1, wherein the antisense oligonucleotide is complementary to a region of TRPM-2 mRNA including the translation initiation or termination site.
3. The method of claim 2, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 4.
4. The method of claim 2, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 5.
5. The method of claim 2, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 12.
6. A method for treating prostate cancer in an individual suffering from prostate cancer, comprising the steps of initiating androgen-withdrawal to induce apoptotic cell death of prostatic tumor cells in the individual, and administering to the individual a composition effective to inhibit expression of TRPM-2 by the tumor cells, thereby delaying the progression of prostatic tumor cells to an androgen-independent state in an individual.
7. The method of claim 6, wherein the composition effective to inhibit expression of TRPM-2 is an antisense oligonucleotide.
8. The method of claim 7, wherein the antisense oligonucleotide is complementary to a region of TRPM-2 mRNA including the translation initiation or termination site.

- 16 -

9. The method of claim 8, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 4.
10. The method of claim 8, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 5.
11. The method of claim 8, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 12.
12. The method of any of claims 8-11, further comprising the step of administering to the individual a chemotherapy agent.
13. The method of claims 12, wherein the chemotherapy agent is a taxane or mitoxanthrone.
14. The method of any of claims 8-11, further comprising the step of administering to the individual a second antisense oligodeoxynucleotide which inhibits expression of an anti-apoptotic protein other than TRPM-2.
15. The method of claim 14, wherein the second antisense oligodeoxynucleotide is antisense Bcl-2 oligodeoxynucleotide.
16. The method of claim 14, further comprising the step of administering to the individual a chemotherapy agent.
17. The method of claims 16, wherein the chemotherapy agent is a taxane or mitoxanthrone.
18. A method for enhancing the chemo- or radiation sensitivity of cancer cells in an individual suffering from a cancer that expresses TRPM-2 in amounts different from normal tissue of

- 17 -

the same type, comprising administering to the individual a composition effective to inhibit expression of TRPM-2 by cancer cells.

19. The method of claim 12, wherein the composition effective to inhibit expression of TRPM-2 is an antisense oligonucleotide.
20. The method of claim 13, wherein the antisense oligonucleotide is complementary to a region of TRPM-2 mRNA including the translation initiation or termination site.
21. The method of claim 14, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 4.
22. The method of claim 14, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 5.
23. The method of claim 14, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 12.

1/10




FIG. 1

2/10




FIG. 2

3/10




FIG. 3




FIG. 4





FIG. 5

5/10



- ANTISENSE TRPM-2 ODN + ANTISENSE BCL-2 ODN + TAXOL
- ANTISENSE TRPM-2 ODN + MISMATCH BCL-2 ODN + TAXOL
- MISMATCH TRPM-2 ODN + ANTISENSE BCL-2 ODN + TAXOL
- MISMATCH TRPM-2 ODN + MISMATCH BCL-2 ODN + TAXOL

FIG. 6

6/10




FIG. 7A






FIG. 7B

TRPM-2 EXPRESSION
OF PC-3 CELLS
5 DAYS AFTER
RADIATION
(DENSITOMETRY OF
NORTHERN BLOT)

**FIG. 8****FIG. 10**




FIG. 9A




FIG. 9B

9/10




FIG. 11A




FIG. 11B

SUBSTITUTE SHEET (RULE 26)




FIG. 12A




FIG. 12B

SEQUENCE LISTING

<110> Gleave, Martin
Rennie, Paul S.
Miyake, Hideaki
Nelson, Colleen

<120> TRPM-2 ANTISENSE THERAPY

<130> UBC.P-020-WO

<140>

<141>

<150> 60/121,726

<151> 1999-02-26

<160> 14

<170> PatentIn Ver. 2.1

<210> 1

<211> 21

<212> DNA

<213> Murine

<220>

<223> antisense TRPM-2 ODN

<400> 1

gcacagcagg agaattcttca t

21

<210> 2

<211> 21

<212> DNA

<213> Murine

<220>

<223> mismatch control

<400> 2

gcacagcagc aggatcttca t

21

<210> 3

<211> 21

<212> DNA

<213> HUMAN

<220>

<223> antisense TRPM-2 ODN

<400> 3

21

tggagtctttt gcacgcctcg g

<210> 4

<211> 21

<212> DNA

<213> HUMAN

<220>

<223> antisense TRPM-2 ODN

<400> 4

21

cagcagcaga gtcttcatca t

<210> 5

<211> 21

<212> DNA

<213> HUMAN

<220>

<223> antisense TRPM-2 ODN

<400> 5

21

attgtctgag accgtctggt c

<210> 6

<211> 21

<212> DNA

<213> HUMAN

<220>

<223> antisense TRPM-2 ODN

<400> 6

21

ccttcagctt tgtctctgat t

<210> 7

<211> 21

<212> DNA

<213> HUMAN

<220>

<223> antisense TRPM-2 ODN

<400> 7

agcagggagt cgatgcggtc a

21

<210> 8

<211> 21

<212> DNA

<213> HUMAN

<220>

<223> antisense TRPM-2 ODN

<400> 8

atcaagctgc ggacgatgcg g

21

<210> 9

<211> 21

<212> DNA

<213> HUMAN

<220>

<223> antisense TRPM-2 ODN

<400> 9

gcaggcagcc cgtggagttg t

21

<210> 10

<211> 21

<212> DNA

<213> HUMAN

<220>

<223> antisense TRPM-2 ODN

<400> 10

ttcagctgct ccagcaagga g

21

<210> 11

<211> 21

<212> DNA

<213> HUMAN

<220>

<223> antisense TRPM-2 ODN

<400> 11

aatttagggg tcttcctgga g

21

<210> 12

<211> 21

<212> DNA

<213> HUMAN

<220>

<223> antisense TRPM-2 ODN

<400> 12

gctgggcgga gttgggggcc t

21

<210> 13

<211> 18

<212> DNA

<213> Murine

<220>

<223> antisense Bcl-2 ODN

<400> 13

tctcccggct tgcgccat

18

<210> 14

<211> 18

<212> DNA

<213> Murine

<220>

<223> mismatch Bcl-2 ODN

<400> 14

tctcccggca tgggtgcat

18

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/04875

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : CO7H 21/04; A61K 48/00; C12N 15/85; C12Q 1/68

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 91.1, 91.3, 325, 375; 536/23.1, 23.2, 24.5, 24.3, 24.31, 24.33; 514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN, MEDLINE, BIOSIS, CAPLUS, LIFESCI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	SENSIBAR ET AL. "Prevention of Cell Death Induced by Tumor Necrosis Factor alpha in LNCaP Cells by Overexpression of Sulfated Glycoprotein-2 (Clusterin)," Cancer Research. 01 June 1995, Vol. 55, pages 2431-2437, see entire document.	1, 2 ----- 3-23
Y	MILNER ET AL "Selecting effective antisense reagents on combinatorial oligonucleotide arrays," Nature Biotechnology. June 1997, Vol. 15, pages 537-541, see entire document.	1, 2, 6-8, 18-20
Y	BUTTYAN ET AL "Induction of the TRPM-2 Gene in Cells Undergoing Programmed Death. Molecular and Cellular Biology," August 1989, Vol. 9, No. 8, pages 3473-3481, see entire document.	1, 2, 6-8, 18-20



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 30 MAY 2000	Date of mailing of the international search report 11 AUG 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer <i>My Meas</i> KAREN A. LACOURCIERE Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/04875

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MILLAR ET AL. "Localization of mRNAs by in-situ hybridization to the residual body at stages IX-X of the cycle of the rat seminiferous epithelium: fact or artefact? International Journal of Andrology," June 1994, Vol. 17, pages 149-160.	1-23
A	DARBY ET AL. "Vascular Expression of Clusterin in Experimental Cyclosporine Nephrotoxicity. Experimental Nephrology," July 1995, Vol. 3, No. 4, pages 234-239.	1-23

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/04875

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/6, 91.1, 91.3, 325, 375; 536/23.1, 23.2, 24.5, 24.3, 24.31, 24.33; 514/44

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 August 2000 (31.08.2000)

PCT

(10) International Publication Number
WO 00/49937 A3

- (51) International Patent Classification⁷: **C07H 21/04**,
A61K 48/00, C12N 15/85, C12Q 1/68
- (21) International Application Number: PCT/US00/04875
- (22) International Filing Date: 25 February 2000 (25.02.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/121,726 26 February 1999 (26.02.1999) US
- (71) Applicant (for all designated States except US): **THE UNIVERSITY OF BRITISH COLUMBIA** [CA/CA];
2075 Westbrook Mall, Vancouver, British Columbia V6T 1Z1 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **GLEAVE, Martin** [CA/CA]; 4693 Drummond Drive, Vancouver, British Columbia V6R 1E8 (CA). **RENNIE, Paul, S.** [CA/CA]; 3731 Ellsmore Avenue, Richmond, British Columbia V7C 1S3 (CA). **MIYAKE, Hideaki** [JP/CA]; 2638 Ash Street, #204, Vancouver, British Columbia V5Z 4K3 (CA). **NELSON, Colleen** [US/CA]; 3299, 137A Street, Surrey, British Columbia V4P 2B5 (CA).
- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— With international search report.
- (88) Date of publication of the international search report:
7 December 2000
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TRPM-2 ANTISENSE THERAPY

WO 00/49937 A3

(57) Abstract: It has now been determined that antisense therapy which reduces the expression of TRPM-2 provides therapeutic benefits in the treatment of cancer. In particular, such antisense therapy can be applied in treatment of prostate cancer and renal cell cancer. Addition of antisense TRPM-2 ODN to prostatic tumor cells *in vivo* is effective for delaying the onset of androgen independence. Thus, prostate cancer can be treated in an individual suffering from prostate cancer by initiating androgen-withdrawal to induce apoptotic cell death of prostatic tumor cells in the individual, and administering to the individual a composition effective to inhibit expression of TRPM-2 by the tumor cells, thereby delaying the progression of prostatic tumor cells to an androgen-independent state in an individual. Combined use of antisense TRPM-2 and taxanes synergistically enhances cytotoxic chemosensitivity of androgen-independent prostate cancer. In addition, it has also been found that antisense TRPM-2 has beneficial effect for other cancer types. Specifically, antisense TRPM-2 ODN enhances chemosensitivity in human Renal cell cancer, a normally chemoresistant disease with no active chemotherapeutic agent having an objective response rate higher than 10 %. Radiation sensitivity is also enhanced when cells expressing TRPM-2 are treated with antisense TRPM-2 ODN. Thus, the antisense TRPM-2 ODNs can be used to enhance hormone sensitivity, chemosensitivity and radiation sensitivity of a variety of cancer types in which expression of TRPM-2 has been observed.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/04875

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : CO7H 21/04; A61K 48/00; C12N 15/85; C12Q 1/68

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 91.1, 91.3, 325, 375; 536/23.1, 23.2, 24.5, 24.3, 24.31, 24.33; 514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN, MEDLINE, BIOSIS, CAPLUS, LIFESCI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	SENSIBAR ET AL. "Prevention of Cell Death Induced by Tumor Necrosis Factor alpha in LNCaP Cells by Overexpression of Sulfated Glycoprotein-2 (Clusterin)," Cancer Research. 01 June 1995, Vol. 55, pages 2431-2437, see entire document.	1, 2 ----- 3-23
Y	MILNER ET AL "Selecting effective antisense reagents on combinatorial oligonucleotide arrays," Nature Biotechnology. June 1997, Vol. 15, pages 537-541, see entire document.	1, 2, 6-8, 18-20
Y	BUTTYAN ET AL "Induction of the TRPM-2 Gene in Cells Undergoing Programmed Death. Molecular and Cellular Biology," August 1989, Vol. 9, No. 8, pages 3473-3481, see entire document.	1, 2, 6,-8, 18-20

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

30 MAY 2000

Date of mailing of the international search report

11 AUG 2000

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

KAREN A. LACOURCIERE

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/04875

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MILLAR ET AL. "Localization of mRNAs by in-situ hybridization to the residual body at stages IX-X of the cycle of the rat seminiferous epithelium: fact or artefact? International Journal of Andrology," June 1994, Vol. 17, pages 149-160.	1-23
A	DARBY ET AL. "Vascular Expression of Clusterin in Experimental Cyclosporine Nephrotoxicity. Experimental Nephrology," July 1995, Vol. 3, No. 4, pages 234-239.	1-23

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/04875

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/6, 91.1, 91.3, 325, 375; 536/23.1, 23.2, 24.5, 24.3, 24.31, 24.33; 514/44

- 15 -

CLAIMS

1. A method for delaying progression of prostatic tumor cells to an androgen- independent state, comprising treating androgen-sensitive prostatic tumor cells with an antisense oligonucleotide which inhibits expression of TRPM-2 by the tumor cells.
2. The method of claim 1, wherein the antisense oligonucleotide is complementary to a region of TRPM-2 mRNA including the translation initiation or termination site.
3. The method of claim 2, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 4.
4. The method of claim 2, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 5.
5. The method of claim 2, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 12.
6. A method for treating prostate cancer in an individual suffering from prostate cancer, comprising the steps of initiating androgen-withdrawal to induce apoptotic cell death of prostatic tumor cells in the individual, and administering to the individual a composition effective to inhibit expression of TRPM-2 by the tumor cells, thereby delaying the progression of prostatic tumor cells to an androgen-independent state in an individual.
7. The method of claim 6, wherein the composition effective to inhibit expression of TRPM-2 is an antisense oligonucleotide.
8. The method of claim 7, wherein the antisense oligonucleotide is complementary to a region of TRPM-2 mRNA including the translation initiation or termination site.

- 16 -

9. The method of claim 8, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 4.
10. The method of claim 8, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 5.
11. The method of claim 8, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 12.
12. The method of any of claims 8-11, further comprising the step of administering to the individual a chemotherapy agent.
13. The method of claims 12, wherein the chemotherapy agent is a taxane or mitoxanthrone.
14. The method of any of claims 8-11, further comprising the step of administering to the individual a second antisense oligodeoxynucleotide which inhibits expression of an anti-apoptotic protein other than TRPM-2.
15. The method of claim 14, wherein the second antisense oligodeoxynucleotide is antisense Bcl-2 oligodeoxynucleotide.
16. The method of claim 14, further comprising the step of administering to the individual a chemotherapy agent.
17. The method of claims 16, wherein the chemotherapy agent is a taxane or mitoxanthrone.
18. A method for enhancing the chemo- or radiation sensitivity of cancer cells in an individual suffering from a cancer that expresses TRPM-2 in amounts different from normal tissue of

- 17 -

the same type, comprising administering to the individual a composition effective to inhibit expression of TRPM-2 by cancer cells.

19. The method of claim 12, wherein the composition effective to inhibit expression of TRPM-2 is an antisense oligonucleotide.
20. The method of claim 13, wherein the antisense oligonucleotide is complementary to a region of TRPM-2 mRNA including the translation initiation or termination site.
21. The method of claim 14, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 4.
22. The method of claim 14, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 5.
23. The method of claim 14, wherein the antisense oligonucleotide has the sequence given by SEQ ID No. 12.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



Spec!

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61B		A2	(11) International Publication Number: WO 00/49937
			(43) International Publication Date: 31 August 2000 (31.08.00)
(21) International Application Number: PCT/US00/04875 (22) International Filing Date: 25 February 2000 (25.02.00) (30) Priority Data: 60/121,726 26 February 1999 (26.02.99) US (71) Applicant (for all designated States except US): THE UNIVERSITY OF BRITISH COLUMBIA [CA/CA]; 2075 Westbrook Mall, Vancouver, British Columbia V6T 1Z1 (CA). (72) Inventors; and (75) Inventors/Applicants (for US only): <u>GLEAVE</u> , Martin [CA/CA]; 4693 Drummond Drive, Vancouver, British Columbia V6R 1E8 (CA). <u>RENNIE</u> , Paul, S. [CA/CA]; 3731 Ellsmore Avenue, Richmond, British Columbia V7C 1S3 (CA). <u>MIYAKE</u> , Hideaki [JP/CA]; 2638 Ash Street, #204, Vancouver, British Columbia V5Z 4K3 (CA). <u>NELSON</u> , Colleen [US/CA]; 3299, 137A Street, Surrey, British Columbia V4P 2B5 (CA). (74) Agents: PARSONS, Nancy, J. et al.; Oppedahl & Larson LLP, 611 Main Street, P.O. Box 5270, Frisco, CO 80443-5270 (US).			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: TRPM-2 ANTISENSE THERAPY			
(57) Abstract <p>It has now been determined that antisense therapy which reduces the expression of TRPM-2 provides therapeutic benefits in the treatment of cancer. In particular, such antisense therapy can be applied in treatment of prostate cancer and renal cell cancer. Addition of antisense TRPM-2 ODN to prostatic tumor cells <i>in vivo</i> is effective for delaying the onset of androgen independence. Thus, prostate cancer can be treated in an individual suffering from prostate cancer by initiating androgen-withdrawal to induce apoptotic cell death of prostatic tumor cells in the individual, and administering to the individual a composition effective to inhibit expression of TRPM-2 by the tumor cells, thereby delaying the progression of prostatic tumor cells to an androgen-independent state in an individual. Combined use of antisense TRPM-2 and taxanes synergistically enhances cytotoxic chemosensitivity of androgen-independent prostate cancer. In addition, it has also been found that antisense TRPM-2 has beneficial effect for other cancer types. Specifically, antisense TRPM-2 ODN enhances chemosensitivity in human Renal cell cancer, a normally chemoresistant disease with no active chemotherapeutic agent having an objective response rate higher than 10 %. Radiation sensitivity is also enhanced when cells expressing TRPM-2 are treated with antisense TRPM-2 ODN. Thus, the antisense TRPM-2 ODNs can be used to enhance hormone sensitivity, chemosensitivity and radiation sensitivity of a variety of cancer types in which expression of TRPM-2 has been observed.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

161
REC 22 MAY 2001

WFO PCT

Applicant's or agent's file reference UBCP020WO	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/US00/04875	International filing date (day/month/year) 25 FEBRUARY 2000	Priority date (day/month/year) 26 FEBRUARY 1999
International Patent Classification (IPC) or national classification and IPC Please See Supplemental Sheet.		
Applicant THE UNIVERSITY OF BRITISH COLUMBIA		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.


2. This REPORT consists of a total of 4 sheets.

☒ This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority. (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of 4 sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of report with regard to novelty, inventive step or industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand 13 SEPTEMBER 2000	Date of completion of this report 24 APRIL 2001
Name and mailing address of the IPEA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer  KAREN A. LACOURCIERE
Facsimile No. (703) 305-3230	Telephone No. (703) 308-0196

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US00/04875

I. Basis of the report

1. With regard to the elements of the international application:*

☐ the international application as originally filed☒ the description:

pages (See Attached) _____, as originally filed
pages _____, filed with the demand
pages _____, filed with the letter of _____

☒ the claims:

pages (See Attached) _____, as originally filed
pages _____, as amended (together with any statement) under Article 19
pages _____, filed with the demand
pages _____, filed with the letter of _____

☒ the drawings:

pages (See Attached) _____, as originally filed
pages _____, filed with the demand
pages _____, filed with the letter of _____

☒ the sequence listing part of the description:

pages (See Attached) _____, as originally filed
pages _____, filed with the demand
pages _____, filed with the letter of _____

2. With regard to the language, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language _____ which is:

☐ the language of a translation furnished for the purposes of international search (under Rule 23.1(b)).☐ the language of publication of the international application (under Rule 48.3(b)).☐ the language of the translation furnished for the purposes of international preliminary examination (under Rules 55.2 and/or 55.3).

3. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

☒ contained in the international application in printed form.☒ filed together with the international application in computer readable form.☐ furnished subsequently to this Authority in written form.☐ furnished subsequently to this Authority in computer readable form.☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.4. ☒ The amendments have resulted in the cancellation of:☒ the description, pages NONE☒ the claims, Nos. NONE☒ the drawings, sheets/fig. NONE5. ☐ This report has been drawn as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).**

* Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17).

**Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US00/04875

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**1. statement**

Novelty (N)	Claims <u>1-28</u>	YES
	Claims <u>NONE</u>	NO
Inventive Step (IS)	Claims <u>1-28</u>	YES
	Claims <u>NONE</u>	NO
Industrial Applicability (IA)	Claims <u>1-28</u>	YES
	Claims <u>NONE</u>	NO

2. citations and explanations (Rule 70.7)

Claims 1-28 meet the criteria set out in PCT Article 33(2)-(4), because the prior art does not teach or fairly suggest methods of treatment for prostate cancer in an individual using a composition which inhibits expression of TRPM-2, including antisense oligonucleotides nor does it teach or fairly suggest the method of delaying progression of prostatic tumor cells to an androgen independent state in vivo, as claimed. Further, the prior art does not teach or fairly suggest methods of enhancing the chemo- or radiation sensitivity of cancer cells in a individual using a composition which inhibits expression of TRPM-2, including antisense oligonucleotides. The prior art does not teach or fairly suggest methods of delaying progression of prostatic cancer cells to an androgen insensitive state, in vivo or in vitro, using an antisense oligonucleotide comprising SEQ ID NO:4, 5, or 12.

----- NEW CITATIONS -----

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US00/04875

Supplemental Box

(To be used when the space in any of the preceding boxes is not sufficient)

Continuation of: Boxes I - VIII

Sheet 10

CLASSIFICATION:

The International Patent Classification (IPC) and/or the National classification are as listed below:

IPC(7): CO7H 21/04; A61K 48/00; C12N 15/85; C12Q 1/68 and US Cl.: 435/6, 91.1, 91.3, 325, 375; 536/23.1, 23.2, 24.5, 24.3, 24.31, 24.33; 514/44

I. BASIS OF REPORT:

This report has been drawn on the basis of the description,

page(s) 1-14, as originally filed.

page(s) NONE, filed with the demand.

and additional amendments:

NONE

This report has been drawn on the basis of the claims,

page(s) NONE, as originally filed.

page(s) NONE, as amended under Article 19.

page(s) NONE, filed with the demand.

and additional amendments:

Pages 15-~~18~~, filed with the letter of 20 March 2001.

This report has been drawn on the basis of the drawings,

page(s) 1-15, as originally filed.

page(s) NONE, filed with the demand.

and additional amendments:

NONE

This report has been drawn on the basis of the sequence listing part of the description:

page(s) 1-4, as originally filed.

pages(s) NONE, filed with the demand.

and additional amendments:

NONE

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Commissioner
US Department of Commerce
United States Patent and Trademark
Office, PCT
2011 South Clark Place Room
CP2/5C24
Arlington, VA 22202
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 23 January 2001 (23.01.01)	
International application No. PCT/US00/04875	Applicant's or agent's file reference UBCP020WO
International filing date (day/month/year) 25 February 2000 (25.02.00)	Priority date (day/month/year) 26 February 1999 (26.02.99)
Applicant GLEAVE, Martin et al	

1. The designated Office is hereby notified of its election made:

☒

in the demand filed with the International Preliminary Examining Authority on:

13 September 2000 (13.09.00)

☐

in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was

☐

was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer Kiwa Mpay Telephone No.: (41-22) 338.83.38
--	--

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF THE RECORDING OF A CHANGE

(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

From the INTERNATIONAL BUREAU

To:

PARSONS, Nancy, J.
Oppedahl & Larson LLP
611 Main Street
P.O. Box 5270
Frisco, CO 80443-5270
ETATS-UNIS D'AMERIQUE

Date of mailing (day/month/year) 07 February 2001 (07.02.01)	
Applicant's or agent's file reference UBCP020WO	IMPORTANT NOTIFICATION
International application No. PCT/US00/04875	International filing date (day/month/year) 25 February 2000 (25.02.00)

1. The following indications appeared on record concerning: <input checked="" type="checkbox"/> the applicant <input checked="" type="checkbox"/> the inventor <input type="checkbox"/> the agent <input type="checkbox"/> the common representative		
Name and Address RENNIE, Paul, S. 3731 Ellsmore Avenue Richmond, British Columbia V7C 1S3 Canada	State of Nationality CA	State of Residence CA
	Telephone No.	
	Facsimile No.	
	Teleprinter No.	
2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning: <input type="checkbox"/> the person <input type="checkbox"/> the name <input checked="" type="checkbox"/> the address <input type="checkbox"/> the nationality <input type="checkbox"/> the residence		
Name and Address RENNIE, Paul, S. 3731 Ullsmore Avenue Richmond, British Columbia V7C 1S3 Canada	State of Nationality CA	State of Residence CA
	Telephone No.	
	Facsimile No.	
	Teleprinter No.	
3. Further observations, if necessary:		
4. A copy of this notification has been sent to: <div style="display: flex; justify-content: space-between;"> <div> <input checked="" type="checkbox"/> the receiving Office <input type="checkbox"/> the International Searching Authority <input checked="" type="checkbox"/> the International Preliminary Examining Authority </div> <div> <input type="checkbox"/> the designated Offices concerned <input checked="" type="checkbox"/> the elected Offices concerned <input type="checkbox"/> other: </div> </div>		

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer Kari Huynh-Khuong Telephone No.: (41-22) 338.83.38
--	--